



Status
 ● Presence
 ● Breeding season
 ● Month with eggs

recorded from
 80 squares
 23%

Ostrich

Struthio camelus
 424 plots from 2,344 records

Virgin Yellow-vented Bulbul squares **9**
 Single YvB month **16**
 Two YvB months **24**

| | Seasonality | | |
|-----------|-------------|----|------|
| | presence | bs | eggs |
| January | 270 | 1 | 5 |
| February | 235 | 1 | 1 |
| March | 172 | 1 | 1 |
| April | 152 | - | 2 |
| May | 156 | - | - |
| June | 222 | 7 | - |
| July | 187 | 14 | 4 |
| August | 184 | 18 | 6 |
| September | 130 | 12 | 9 |
| October | 156 | 19 | 11 |
| November | 129 | 15 | 6 |
| December | 206 | 7 | 4 |
| month 13 | 1 | | |

from Britton (1980).
 A locally common resident of dry open and wooded grassland, mainly below 2000m. The nominate race reaches Karamoja in NE Uganda, ranging south to Debasien GR. The Somali race *molybdophanes* occurs in Kenya south to Laikipia, Isiola, Meru NP and Tsavo East NP. The race *massaicus* occurs north of the Kenya highlands at Timau and Laikipia, and ranges from Sotik, Nairobi and Tsavo West NP south through the dry interior of Tanzania to the Katavi Plain and Ruaha NP.



Tanzania Bird Atlas
 Preliminary Map
 August 2012
 From 984,844
 Database records

| | | | |
|---|---|---|---|
| J | F | M | A |
| M | J | J | A |
| S | O | N | D |
| # | # | # | # |

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E