

Zebra Waxbill

Amandava subflava

319 plots from 596 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality

	presence	bs	eggs
January	85	-	-
February	38	-	-
March	31	-	-
April	30	2	-
May	40	1	1
June	50	8	3
July	40	3	3
August	44	1	3
September	37	1	-
October	53	-	-
November	60	-	-
December	61	-	-
month 13	1		

This is nowhere an easy bird and the only locality for which we have records in each month is Peramiho near Songea.


recorded from 109 squares 31 %

Status

- Presence
- Breeding season
- Month with eggs

Note the neatly defined breeding season. The Zanzibar data is from Pakenham 1979 but be aware that he left Zanzibar in 1956 so much will have changed since then.

from Britton (1980)
 A widespread but seldom common resident and wanderer frequenting long grass, swamp edges and cultivation up to 2200m seldom far from permanent water and usually in high rainfall areas. The nominate race ranges widely in Uganda north to Masindi, Kabalega Falls NP, Teso and Kidepo Valley NP, and in Siaya, Kisumu and Uasin Gishu in W Kenya. The race *clarkei* is inexplicably local, occurring in the coastal lowlands at Mikindani, Soga, Zanzibar, Pemba, Mafia and Witu; the E edge of the Kenya highlands from Ngong and Nairobi to Mt Kenya; interior E TZ from Arusha NP, the Crater Highlands and the foothills of the Ulugurus to Njombe and Songea and SW TZ from the Ufipa Plateau to Mpanda. Sight records from West Lake, Kibondo and N Tabora in NW Tanzania cannot be assigned racially.



Tanzania Bird Atlas
 Preliminary Map
 December 2015
 From 1,144,740
 Database records

J F M A
 M J J A
 S O N D
 # # # #

symbols give month of record within square, not the locality within square