



In the SE near Songea there is a small population of what appears to be *angolensis* but with a very small red cheek patch.

recorded from 223 squares 64%

**Status**  
 ● Presence  
 ● Breeding season  
 ● Month with eggs

# Red-cheeked Cordonbleu

*Uraeginthus bengalus*

1,242 plots from 4,157 records

Virgin Yellow-vented Bulbul squares **9**  
 Single YvB month **16**  
 Two YvB months **24**

	Seasonality		
	presence	bs	eggs
January	458	6	4
February	379	7	6
March	329	9	3
April	219	12	9
May	247	8	6
June	326	10	3
July	363	5	1
August	378	4	3
September	294	2	1
October	353	9	1
November	317	3	-
December	366	2	-
month 13	4		



**Tanzania Bird Atlas**  
 Preliminary Map  
 August 2014  
 From 1,072,733  
 Database records

J F M A  
 M J J A  
 S O N D  
 # # # #  
 symbols give month of record within square, not the locality within square

from Britton (1980).  
 A widespread and usually common resident from sea-level to at least 2300 m, typically in higher rainfall areas or moist habitats in drier country, though it is sometimes common in arid and semi-arid country in NE Kenya. Four races occur: *littoralis* in coastal and SE Kenya inland to the Tsavo region, and NE Tanzania from Tanga to the North Pare Mts; *brunneigularis* in the C Kenya highlands and some plateau areas east of the rift, north and east to Moyale and Wajir, intergrading with the coastal race in the southeast; nominate in Kenya west of *brunneigularis* (not Turkana) and throughout Uganda; and *ugogoensis* in the remainder of Tanzania south and east to Rukwa, the Usangu Flats, N Njombe, the Iringa area, Kingolwira, Handeni, the Arusha-Moshi area and Mt Kilimanjaro.

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E