

note the "new" name in general use is *Crithagra dorsostriata*

White-bellied Canary

Serinus dorsostriatus

518 plots from 1,462 records


Status
 ● Presence
 ● Breeding season
 ● Month with eggs

Occurs in 110 squares 31%

Virgin Yellow-vented Bulbul squares **9**
 Single YvB month **16**
 Two YvB months **24**

| | Seasonality | | |
|-----------|-------------|----|------|
| | presence | bs | eggs |
| January | 139 | 3 | 1 |
| February | 121 | 1 | 1 |
| March | 125 | 1 | - |
| April | 101 | 2 | - |
| May | 102 | 1 | 3 |
| June | 139 | 1 | 1 |
| July | 142 | 1 | - |
| August | 145 | 2 | 2 |
| September | 97 | 1 | 1 |
| October | 104 | 1 | - |
| November | 80 | 1 | - |
| December | 139 | 3 | 1 |

from Britton (1980).
 A widely distributed and generally common resident and wanderer in bushed and wooded grassland, bushland, woodland and cultivation in low and medium rainfall areas, most typically in arid and semi-arid country at 400-1400 m, but as high as 1800 m in Narok and 2650m at Mt Nyiru. It occurs in E Karamoja in NE Uganda, and widely in plateau country in Kenya and N Tanzania south to about 6°S in Dodoma and Singida, including NE Tabora, Shinyanga, Mwanza, Biharamulo, the Sibiti River and Serengeti NP in NW Tanzania, and Narok and the drier parts of Nyanza in SW Kenya. The southern nominate race and the northern maculicollis intergrade over a considerable area in the vicinity of the equator in E Kenya. Ripley & Heinrich (1966) have, however, extended the range of maculicollis south to Same and the base of Mt Meru in extreme NE Tanzania.



Tanzania Bird Atlas
 Preliminary Map
 March 2013
 From 1,026,411
 Database records

| | | | |
|---|---|---|---|
| J | F | M | A |
| M | J | J | A |
| S | O | N | D |
| # | # | # | # |

symbols give month of record within square, not the locality within square