

# Stripe-breasted Seedeater

*Crithagra reichardi*

155 plots from 257 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

	Seasonality		
	presence	bs	eggs
January	23	-	-
February	12	-	-
March	23	-	-
April	20	-	-
May	20	1	-
June	17	-	-
July	31	-	-
August	28	-	-
September	24	-	-
October	26	-	-
November	19	-	-
December	12	-	-
month 13	1	-	-

**SO FEW BREEDING SEASON RECORDS !!!!**

- Status**
- Presence
  - Breeding season
  - Month with eggs

These 2 populations may well be genuine isolated outliers.

from Britton (1980).  
 The nominate race is endemic to miombo woodland and associated cultivation. In Tanzania it is a rather local but often common resident at 500-1500 m, from Masasi, Songea and Ufipa north to N Iringa, C Tabora and Kibondo. The race *striatipectus* is locally distributed and uncommon in lightly wooded country in the highlands of W and C Kenya at 1600-1800 m, from Limuru to Mt Ololokwe, Mt Nyiru, the Tugen Hills, Eldama Ravine, Kabarnet, the Elgeyu Escarpment and Mt Elgon, extending to the Lake Victoria basin at Muhoroni.



**Tanzania Bird Atlas**  
 Preliminary Map  
 March 2013  
 From 1,031,269  
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

