

Long-toed Plover

Vanellus crassirostris

248 plots from 550 records

- Status**
- Presence
 - Breeding season
 - Month with eggs

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Occurs in 78 squares 22%

	Seasonality		
	presence	bs	eggs
January	107	-	1
February	60	1	-
March	46	-	-
April	16	-	-
May	13	-	-
June	25	-	1
July	45	-	2
August	47	1	-
September	40	2	-
October	62	1	-
November	47	-	1
December	25	3	-
month 13	2		

from Britton (1980).
 The nominate race is a locally common resident from sea level to 2200m in permanent swamps and swampy edges of lakes and rivers with abundant floating vegetation. It occurs throughout Uganda except the extreme northeast, but has a curiously fragmented distribution in Kenya and Tanzania. In W Kenya it is common in the Lake Victoria basin, and locally in the peripheral highlands in Uasin Gishu, Trans Nzoia and Nandi, but to the east it occurs only at Lake Naivasha, Amboseli, Lake Jipe and the lower Tana (including Karawa). In Tanzania it is also a bird of the west, where it is locally distributed east to Tarangire NP and the Kilombero River, south to the Ugalla River and Rukwa.

There is a suggestion here of an eastwards expansion in recent years. Note that these recent records from Bagamoyo are not yet in our database.



Tanzania Bird Atlas
 Preliminary Map
 June 2012
 From 965,876
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

