

# Red-billed Hornbill

*Tockus erythrorhynchus*

238 plots from 795 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **25**

Two YvB months **25**

	Seasonality		
	presence	bs	eggs
January	88	3	1
February	65	3	-
March	51	2	-
April	59	1	-
May	43	1	-
June	51	-	-
July	83	-	1
August	85	-	-
September	56	-	-
October	71	-	-
November	45	-	1
December	81	3	1

## Status

- Presence
- Breeding season
- Month with eggs

## Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900

To accurately define the range of this species, especially with reference to Ruaha Hornbill we need GIS data from all western and SW border squares. The suspicion is that this species only occurs east of the Ruft Valley wall, east of the Mbulu Highlands and north of the Kondoa-Kibaya Ridge. There are a few records to clarify on the edge of this range that could be good for either species.

3504A Dongobesh-Haydom Dave Peterson  
 Katesh Paul Oliver  
 3504C Katesh Marc Baker  
 3605C Zoissa Bruce Hills  
 3706C Kimamba John Beesley

GPS records from ALL squares marked with GREEN circles PLEASE.

Tanzania Bird Atlas  
 Preliminary Map  
 January 2009  
 From 828,055  
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records within each square

