

Red-fronted Tinkerbird

Pogoniulus pusillus

650 plots from 1,683 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Occurs in 155 squares 44%

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	166	3	-
February	164	2	-
March	169	3	-
April	111	3	2
May	78	4	-
June	107	-	-
July	131	-	-
August	150	2	-
September	137	-	-
October	154	-	-
November	145	3	-
December	144	5	-

from Britton (1980).
The race affinis is common in open woodland and bushland, gardens and occasionally forest edges from sea-level to 2200 m (mainly below 1800 m), typically in drier country than the allopatric *P. chrysoconus* with which it forms a superspecies. Apart from western areas occupied by *chrysoconus* it is wide-ranging in Kenya north to Lokori, the Turkwell River, the Horr Valley, Loiyengalani and Marsabit, and in Tanzania south and west to the Rufiji River, Nachingwea, Ruaha NP, Serengeti and Tabora (including Ufipa). In Uganda it occurs in E Karamoja and much of Teso, while *chrysoconus* reaches Teso at Soroti and Komolo.

Status
● Presence
● Breeding season
● Month with eggs

many of these odd records are likely to be ID errors, if any are yours please let us know !!



Tanzania Bird Atlas
Preliminary Map
May 2012
From 965,876
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

