

Brown-eared Woodpecker

Campethera caroli

8 plots from 16 records

Virgin Yellow-vented Bulbul squares	9
Single YvB month	16
Two YvB months	24

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	-	-	-
February	-	-	-
March	-	-	-
April	-	-	-
May	-	-	-
June	-	-	-
July	4	-	-
August	4	-	-
September	2	-	-
October	2	-	-
November	2	2	-
December	-	-	-

Status
 ● Presence
 ● Breeding season
 ● Month with eggs

Occurs in only 2 squares

Our only records for this bird come from Minziro FR in the extreme NW. It is widespread in suitable habitat in Uganda south of 2 degrees north. The seasonality of the few records simply reflects the months when birders have visited this site.

from Britton (1980). Brown-eared Woodpecker *C. caroli*. A common bird of forest and thick secondary habitats at 700-1800m occurring in S and W Uganda from Budongo to the Impenetrable and Sango Bay forests, east to Kifu, Mabira and Mt Elgon, extending to Kakamega Forest in W Kenya and Bukoba in NW Tanzania.



Tanzania Bird Atlas
 Preliminary Map
 March 2013
 From 1,031,732
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

