



Semi-collared Flycatcher

Ficedula semitorquata

28 plots from 47 records

Virgin Yellow-vented Bulbul squares 7

Single YvB month 14

Two YvB month 20

Status

- Presence
- Breeding season
- Month with eggs

Seasonality

	presence	br. season	eggs
January	11	-	-
February	12	-	-
March	2	-	-
April	1	-	-
May	-	-	-
June	1	-	-
July	-	-	-
August	-	-	-
September	2	-	-
October	1	-	-
November	4	-	-
December	13	-	-



Tanzania
BIRD ATLAS
Tanzania Bird Atlas
Preliminary Map
February 2016
From 1,145,267
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square

from Britton (1980).
Most records of this Palearctic migrant are from miombo woodland in Tanzania (west of 36° E) where it is regular and sometimes reasonably common in January-February in parts of Tabora and Iringa, and fairly common on passage (28 September – 6 Nov, 20 February – 4 March) in Kibondo. Both the nominate race and *semitorquata* occur. Specimens from Kibondo on 24 February and 6 October, and Ng'iya in W Kenya on 2 October, are the only definite records of nominate birds, while there are specimens or good sight records of adult male *semitorquata* from Iringa, Njombe and Tabora in January-February, Kibondo on 21 February, Budongo Forest and Butiaba in W Uganda on 22 February and 15 March, and the Mara River in W Kenya on 3 March.

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E