



African Goshawk

Accipiter tachiro

691 plots from 1,820 records

recorded from
194 squares
55.4 %

- Presence
- Breeding season
- Month with eggs

- Virgin Yellow-vented Bulbul squares 4
- Single YvB month 13
- Two YvB months 18

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	142	4	-
February	149	4	-
March	130	14	1
April	94	1	-
May	106	3	-
June	135	7	-
July	191	10	-
August	166	3	-
September	177	1	-
October	162	4	-
November	170	5	-
December	138	3	-

from Britton 1980.
Widespread in forest, rich woodland and gardens up to 3000 m in the less arid parts of East Africa, including Zanzibar, Pemba and Mafia. It is probably the commonest accipiter in most suitable habitat. Apart from Pemba, where the small richly coloured *pembaensis* is endemic (Benson and Elliott 1975), the race *sparsifasciatus* ranges throughout East Africa.



Tanzania Bird Atlas
Preliminary Map
August 2017
From 1,180,093
Database records

J F M A
M J J A
S O N D

symbols give month of
record within square, not
the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E