

# Long-crested Eagle

*Lophaetus occipitalis*

959 plots from 2,794 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **12**

Two YvB months **21**


### Status

- Presence
- Breeding season
- Month with eggs

### Seasonality

	presence	bs	eggs
January	301	-	-
February	288	-	1
March	230	3	-
April	158	4	-
May	159	1	-
June	201	-	-
July	258	4	1
August	245	1	3
September	215	8	-
October	234	6	1
November	228	1	1
December	237	1	1
month 13	3		

from Britton (1980).  
 Widespread and locally numerous in higher rainfall areas up to 3000 m, though generally uncommon above 2000 m. An adaptable woodland and forest edge species which is especially common in areas partially cleared for agriculture, even when heavily settled. It takes large numbers of rodents and is generally considered beneficial to man. An invasion in Serengeti NP coincided with exceptional numbers of grass rats *Arvicanthis*.



**Tanzania Bird Atlas**  
 Preliminary Map  
 February 2015  
 From 1,113,718  
 Database records

J F M A  
 M J J A  
 S O N D  
 # # # #

symbols give month of record within square, not the locality within square

20°E 20°E 24°E 22°E 22°E 24°E 25°E 26°E 27°E 28°E 29°E 40°E 41°E