



The few records along the shores of our major lakes surely relates to poor coverage. Mahale, Rubondo and Bukoba are examples of what this map should show.

recorded from
88 squares
25 %

Status
● Presence
● Breeding season
● Month with eggs

Osprey

Pandion haliaetus

246 plots from 432 records

Virgin Yellow-vented Bulbul squares **9**
Single YvB month **12**
Two YvB months **21**

	Seasonality		
	presence	bs	eggs
January	102	-	-
February	50	-	-
March	46	-	-
April	27	-	-
May	7	-	-
June	12	-	-
July	23	-	-
August	15	-	-
September	24	-	-
October	37	-	-
November	37	-	-
December	50	-	-
month 13	2	-	-



Tanzania Bird Atlas
Preliminary Map
February 2015
From 1,110,908
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

from Britton (1980).
The nominate race is a reasonably common Palearctic migrant and winter visitor to lakes, rivers and coasts, mainly September-March, but regularly as early as July-August and as late as April. Few oversummer and there is no proper breeding record. It occurs in suitable localities throughout, including Zanzibar and Pemba, but coastal birds are virtually confined to creeks, estuaries and islets.