



There are a few potential errors here in the southern Serengeti and NCA and south into the Wembere where Grey-breasted occurs. That Mkomazi records is also likely to be an error and does this bird really occur in Arusha and 3703C ? All will be checked.

recorded from
255 squares
72.8 %

Red-necked Spurfowl

Pternistis afer

1,213 plots from 4,002 records

- Virgin Yellow-vented Bulbul squares 4
- Single YvB month 13
- Two YvB months 18

Seasonality

presence bs eggs

| Month | presence | bs | eggs |
|-----------|----------|----|------|
| January | 390 | 7 | 1 |
| February | 354 | 7 | 6 |
| March | 369 | 10 | 9 |
| April | 353 | 10 | 12 |
| May | 191 | 10 | 14 |
| June | 284 | 21 | 17 |
| July | 324 | 24 | 20 |
| August | 303 | 14 | 11 |
| September | 270 | 4 | 5 |
| October | 329 | 6 | 2 |
| November | 307 | 3 | 2 |
| December | 308 | 3 | 1 |

from Britton (1980).
Widespread in bushed grassland, wooded grassland, bushland thicket and cultivation from sea-level to 1500 m, except in areas occupied by other bare-throated species. The race *cranchii* is common in SW UG north to Toro, ranging through W TZ south to Ufipa, except around Kasulu where *harterti* occurs. To the east, *cranchii* occurs north of *F. rufopictus* in N Serengeti, the Keekorok-Sand River area of Mara GR and S. Nyanza, and locally on the north shore of L. Victoria east to Fort Ternan. The race *melanogaster* is locally distributed through the coastal lowlands of TZ inland to Korogwe and Songea, hybridizing with *cranchii* from Tukuyu and Njombe to Tabora, Mbulu and Dodoma.



Tanzania Bird Atlas
Preliminary Map
December 2016
From 1,169,150
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E