



## Red-necked Spurfowl

*Pternistis afer*

from 1,205 geo-referenced records

Virgin Yellow-vented Bulbul squares	4
Single YvB month	13
Two YvB months	18

Seasonality	presence	bs	eggs
January	390	7	1
February	354	7	6
March	369	10	9
April	353	10	12
May	191	10	14
June	284	21	17
July	324	24	20
August	303	14	11
September	270	4	5
October	329	6	2
November	307	3	2
December	308	3	1

from Britton (1980). Widespread in bushed grassland, wooded grassland, bushland thicket and cultivation from sea-level to 1500 m, except in areas occupied by other bare-throated species. The race *cranchii* is common in SW UG north to Toro, ranging through W TZ south to Ufipa, except around Kasulu where *harterti* occurs. To the east, *cranchii* occurs north of *F. rufopictus* in N Serengeti, the Keekorok-Sand River area of Mara GR and S. Nyanza, and locally on the north shore of L. Victoria east to Fort Ternan. The race *melanogaster* is locally distributed through the coastal lowlands of TZ inland to Korogwe and Songea, hybridizing with *cranchii* from Tukuyu and Njombe to Tabora, Mbulu and Dodoma.



Tanzania Bird Atlas  
Preliminary Map  
December 2016  
From 1,169,150  
Database records

1° S  
2° S  
3° S  
4° S  
5° S  
6° S  
7° S  
8° S  
9° S  
10° S  
11° S  
12° S

29° E 30° E 31° E 32° E 33° E 34° E 35° E 36° E 37° E 38° E 39° E 40° E 41° E