
Red-necked Spurfowl
Pternistis afer
1,051 geos from 333,422
geo-referenced records
no YvB records 7
Single YvB month 14
Two YvB months 20
from Britton (1980)
Widespread in bushed grassland, wooded grassland bushland thicket and cultivation from sea-level to 1500m, except in areas occupied by other bare-throated species (*F. leucoscepus, F. rufopictus*) with which it is allopatric. The race *cranchii* is very common in SW UG north to Toro, ranging through W TZ south to Ufipa, except around Kasulu where *harterti* occurs. To the east, *cranchii* occurs north of *F. rufopictus* in N Serengeti, the Keekorok-Sand River area of Mara GR and South Nyanza, and locally on the north shore of Lake Victoria east to Fort Ternan. The race *melanogaster* is locally distributed throughout the coastal lowlands of TZ inland to Korogwe, Mahenge and Songea, hybridizing with *cranchii* from Tukuyu and Njombe to Tabora, Mbulu and Dodoma. In the coastal lowlands of KE, the race *leucoparaeus* is common north to the Shimba Hills and north of the Tana river, but it is hardly recoded in between (where *F. leucoscepus* occurs, though only reaching the coast between the Sabaki and Tana rivers).
TANZANIA BIRD ATLAS
Tanzania Bird Atlas
Preliminary Map
October 2015
From 1,139,059
Database records
1°S
2°S
3°S
4°S
5°S
6°S
7°S
8°S
9°S
10°S
11°S
29°E
30°E
31°E
32°E
33°E
34°E
35°E
36°E
37°E
38°E
39°E
40°E
41°E