

Red-necked Spurfowl

Francolinus afer

1,164 plots from 3,396 records


	Seasonality		
	presence	bs	eggs
January	361	5	1
February	306	6	6
March	253	10	6
April	226	10	9
May	172	9	13
June	226	19	16
July	280	22	20
August	282	14	11
September	246	4	5
October	287	5	2
November	273	3	2
December	281	3	1
month 13	1		

Rather many question marks here for us to double check. Does anyone know where this bird meets Grey-breasted ? can you GPS it for us ? Geo map soon.

recorded from 247 squares 71%

Status

- Presence
- Breeding season
- Month with eggs



from Britton (1980). Widespread in bushed grassland, wooded grassland, bushland thicket and cultivation from sea-level to 1500 m, except in areas occupied by other bare-throated species (*F. leucoscepus*, *F. rufopictus*) with which it is allopatric. The race *cranchii* is very common in SW Uganda north to Toro, ranging through W Tanzania south to Ufipa, except around Kasulu where *harterti* occurs. To the east, *cranchii* occurs north of *F. rufopictus* in N Serengeti, the Keekorok-Sand River area of Mara GR and South Nyanza, and locally on the north shore of Lake Victoria east to Fort Ternan. The race *melanogaster* is locally distributed through the coastal lowlands of Tanzania inland to Korogwe, Mahenge and Songea, hybridizing with *cranchii* from Tukuyu and Njombe to Tabora, Mbulu and Dodoma. In the coastal lowlands of Kenya, the race *leucoparaeus* is common north to the Shimba Hills and north of the Tana river, but it is hardly recorded in between (where *F. leucoscepus* occurs, though only reaching the coast between the Sabaki and Tana rivers).



Tanzania Bird Atlas
Preliminary Map
March 2014
From 1,061,639
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square