

Red-necked Spurfowl

Pternistis afer

1,197 plots from 3,839 records


Virgin Yellow-vented Bulbul squares **7**
 Single YvB month **14**
 Two YvB months **20**

Seasonality	presence	bs	eggs
January	374	6	1
February	334	6	6
March	364	10	8
April	346	10	10
May	187	9	14
June	275	20	17
July	308	23	20
August	294	14	11
September	252	4	5
October	305	6	2
November	292	3	2
December	296	3	1

Some of the records enclosed in red may be ID errors for Grey-breasted. That odd record on Pemba is an entry error and has been deleted. Otherwise this is quite a good map for this widespread and locally common species.

recorded from 246 squares 70 %

Status
 ● Presence
 ● Breeding season
 ● Month with eggs



from Britton (1980)
 Widespread in bushed grassland, wooded grassland bushland thicket and cultivation from sea-level to 1500m, except in areas occupied by other bare-throated species (*F. leucoscepus*, *F. rufopictus*) with which it is allopatric. The race *cranchii* is very common in SW UG north to Toro, ranging through W TZ south to Ufipa, except around Kasulu where *harterti* occurs. To the east, *cranchii* occurs north of *F. rufopictus* in N Serengeti, the Keekorok-Sand River area of Mara GR and South Nyanza, and locally on the north shore of Lake Victoria east to Fort Ternan. The race *melanogaster* is locally distributed throughout the coastal lowlands of TZ inland to Korogwe, Mahenge and Songea, hybridizing with *cranchii* from Tukuyu and Njombe to Tabora, Mbulu and Dodoma. In the coastal lowlands of KE, the race *leucoparaeus* is common north to the Shimba Hills and north of the Tana river, but it is hardly recorded in between (where *F. leucoscepus* occurs, though only reaching the coast between the Sabaki and Tana rivers).



Tanzania Bird Atlas
 Preliminary Map
 October 2015
 From 1,139,059
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of record within square, not the locality within square