

Yellow-necked Sparrow

Premistis leucoscepus

371 plots from 1,778 records

Virgin Yellow-vented Bulbul squares **4**

Single YvB month **13**

Two YvB months **18**

recorded from
249 squares
71 %

- Presence
- Breeding season
- Month with eggs

Seasonality
presence by eggs

Month	presence	by	eggs
January	190	1	2
February	174	4	2
March	214	3	2
April	150	3	1
May	98	8	2
June	107	6	8
July	136	7	11
August	153	11	2
September	113	1	-
October	127	1	1
November	107	-	-
December	151	-	-

interesting that we continue to receive the occasional record from the Ruaha ecosystem but it is far from common. Might some birds move south seasonally? Those odd records west of the core range may well be ID errors for Grey-breasted Sparrow L.

from Britton (1980). The race *infuscatus* is common in bushy grassland and cultivation from sea-level to 2400 m, allopatric with other bare-throated species (*F. afer*, *F. rufopectus*). Though wide-ranging in Kenya, it is absent from the humid Lake Victoria basin, Whitehighlands and coastal belt (except between the Sabaki and Tana rivers). It ranges west to Mt Moroto and Kidepo Valley NP in Karamoja, and is locally common on the east shore of Lake Turkana, but hardly occurs in Turkana itself. In Tanzania it occurs in the E plateau south to Dodoma and Morogoro, west to Nduu where it hybridizes with *rufopectus*. Though absent from Serengeti NP it occurs in the Mara GR to the east and south of Keekorok.



Tanzania Bird Atlas
Preliminary Map
December 2016
From 1,173,386
Database records

J F M A
M J J A
S O N D
● ● ● ●

symbols give month of record within square, not the locality within square

