

Yellow-necked Spurfiowl

Pternistis leucoscepus

369 plots from 1,731 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	185	1	2
February	170	4	2
March	212	3	2
April	148	5	1
May	92	8	2
June	102	6	8
July	129	7	11
August	131	11	2
September	108	1	-
October	122	1	1
November	105	-	-
December	149	-	-

recorded from
67 squares
19 %

Status

- Presence
- Breeding season
- Month with eggs

These 3 out of range records and MOST (ALL!!) of the records from the Serengeti will refer to Grey-breasted Spurfiowl.

Photos are required for any claims from the Serengeti NP and especially of any possible hybrids from the Nduu area.




Tanzania Bird Atlas
Preliminary Map
June 2015
From 1,130,669
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

from Britton (1980)
The race *infuscatus* is common in bushy grassland and cultivation from sea-level to 2400 m, allopatric with other bare-throated species (*F. afer*, *F. rufopictus*). Though wide-ranging in Kenya, it is absent from the humid Lake Victoria basin, W highlands and coastal belt (except between the Sabaki and Tana rivers). It ranges west to Mt Moroto and Kidepo Valley NP in Karamoja, and is locally common on the east shore of Lake Turkana, but hardly occurs in Turkana itself. In Tanzania it occurs in the E plateau south to Dodoma and Morogoro, west to Nduu where it hybridizes with *rufopictus*. Though absent from Serengeti NP it occurs in the Mara GR to the east and south of Keekorok.



29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E