

Grey Crowned Crane

Balearica regulorum

524 plots from 1,823 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality

	presence	bs	eggs
January	285	6	3
February	190	7	5
March	169	5	12
April	132	11	8
May	103	2	5
June	114	5	5
July	123	4	2
August	137	1	-
September	93	-	-
October	111	-	-
November	110	3	-
December	161	6	-
month 13	4		

THESE RECORDS ARE FROM FROM 35 YEARS OF DATA COLLECTION. COMPARE WITH THE MAP FOR THE PAST 15 YEARS.

recorded from 115 squares 32 %

Status

- Presence
- Breeding season
- Month with eggs

from Britton (1980)

A widespread resident in pairs and flocks in grassland up to 3000m, breeding in marshy areas but feeding well away from water. Apart from N and E Kenya (north of Maralal and the N Uaso Nyiro River), the coastal lowlands and SE Tanzania, the race *gibbericeps* occurs throughout. It is not known to breed east of Embu, Stony Athi and Arusha NP, and in Tsavo East NP it is only a sporadic visitor. The race *ceciliae* may be only a seasonal visitor from the north, recorded in the Nile Valley south to Pakwach and Kabalega Falls NP and along the shores of Lake Turkana south to Lowarangak and Ileret.

Frame, G.W. 1982. East African Crowned Crane *Balearica regulorum gibbericeps* ecology and behaviour in northern Tanzania. Scopus 6: 60-69.



Tanzania Bird Atlas Preliminary Map July 2015 From 1,135,191 Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

