



Curious that Britton does not mention the southern race *leucoptera*. We have records of this race from Manyara, Katavi and the Usungu but, as far as I recall, no images. Please look out for this white winged race and take some GOOD photos for us. It should be in Tarangire.

recorded from
102 squares
29%

Status
● Presence
● Breeding season
● Month with eggs

Long-toed Plover

Vanellus crassirostris

270 plots from 599 records

Virgin Yellow-vented Bulbul squares **9**
Single YvB month **16**
Two YvB months **24**

	Seasonality		
	presence	bs	eggs
January	114	-	1
February	66	1	-
March	48	-	-
April	16	-	-
May	18	-	-
June	30	-	1
July	49	-	2
August	53	1	-
September	44	2	-
October	66	1	-
November	32	-	-
December	51	-	1
month 13	2		

from Britton (1980).
The nominate race is a locally common resident from sea level to 2200m in permanent swamps and swampy edges of lakes and rivers with abundant floating vegetation. It occurs throughout Uganda except the extreme northeast, but has a curiously fragmented distribution in Kenya and Tanzania. In W Kenya it is common in the Lake Victoria basin, and locally in the peripheral highlands in Uasin Gishu, Trans Nzoia and Nandi, but to the east it occurs only at Lake Naivasha, Amboseli, Lake Jipe and the lower Tana (including Karawa). In Tanzania it is also a bird of the west, where it is locally distributed east to Tarangire NP and the Kilombero River, south to the Ugalla River and Rukwa.



Tanzania Bird Atlas
Preliminary Map
May 2014
From 1,068,711
Database records

J F M A
M J J A
S O N D

symbols give month of
record within square, not
the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E