


Black-winged Stilt

Himantopus himantopus

702 plots from 3,213 records



recorded from
155 squares
44%

- Status**
- Presence
 - Breeding season
 - Month with eggs

- Virgin Yellow-vented Bulbul squares **7**
- Single YvB month **14**
- Two YvB months **20**

Seasonality

	presence	bs	eggs
January	515	2	-
February	309	-	-
March	289	3	2
April	155	-	1
May	138	-	5
June	159	7	13
July	261	6	7
August	245	1	2
September	214	-	-
October	309	1	-
November	258	-	-
December	309	-	-
month 13	2	-	-



Tanzania Bird Atlas
Preliminary Map
April 2015
From 1,125,139
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square

from Britton (1980)
The nominate race is widespread but rather local, favouring the shores of rift valley lakes, coastal salt pans and paddy-fields. It breeds in small numbers at Lake Magadi (annual,) L. Edward, L. Katwe, L. Turkana, Suguta R, Aruba, Ngong Swamps, L. Manyara, Ngare Nairobi and Dar es Salaam. Its occurrence in hundreds at several sites where it seldom or never breeds, mainly between October and April, provides convincing circumstantial evidence that the majority of birds are of Palearctic origin.

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E