



# Collared Pratincole

*Glareola pratincola*

338 plots from 928 records

	Seasonality		
	presence	bs	eggs
January	119	1	-
February	85	-	-
March	71	1	-
April	45	2	2
May	29	-	2
June	43	4	3
July	68	4	11
August	81	3	6
September	107	3	2
October	101	1	1
November	63	-	-
December	70	-	-

from Britton (1980).  
 The race *fuelleborni* is widespread and locally common in wetland and littoral habitats up to 1800 m. All breeding records are from below 1500 m, mainly in low rainfall areas, especially Kabalega Falls NP, Rwenzori NP and Lakes Turkana, Magadi, Manyara and Rukwa. There are few localities where it is present throughout the year. It is a rains breeder in Kenya, Uganda and NE Tanzania, where all egg-laying is between April and September, especially in the wettest months of April and May. Most colonies are on alkaline or sand flats by lakes. Elsewhere in Tanzania, where the usual breeding habitat is dry mudflats left exposed after flood plains have dried out, all egg-laying is in the dry season, between June and September.



Tanzania Bird Atlas  
 Preliminary Map  
 August 2014  
 From 1,072,733  
 Database records

J F M A  
 M J J A  
 S O N D  
 # # # #  
 symbols give month of  
 record within square, not  
 the locality within square