

Senegal Coucal

Centropus senegalensis

115 plots from 256 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

- Status**
- Presence
 - Breeding season
 - Month with eggs

	Seasonality		
	presence	bs	eggs
January	32	2	-
February	19	1	-
March	18	-	-
April	12	-	-
May	9	-	-
June	16	-	-
July	15	-	-
August	28	-	-
September	31	-	-
October	21	-	-
November	19	1	-
December	30	2	-

These records within the Malagarasi catchment may relate to ID errors for Coppery-tailed Coucal or even Bue-headed, esp if they are from wetland habtat. BUT if they are from Borassus flood-plain habitat they may well be Senegal.

from Britton (1980).
 A sometimes common resident of thickets, bushed grassland and forest clearings and edges, less dependent on wet habitats and undergrowth than other coucals. The nominate race is widespread in N Uganda south to Kabalega Falls NP, N Mengo, Lango and Teso, ranging to W Kenya at Kakamega, Mumias and the E slopes of Mt Elgon. There are scattered records of the race *flecki* from interior Tanzania, at Chimala, Ruaha NP, Morogoro and in thickets dominated by Borassus palms on the flood-plain of the Ugalla River.



Tanzania Bird Atlas
 Preliminary Map
 July 2014
 From 1,071,566
 Database records

J F M A
 M J J A
 S O N D
 # # # #

symbols give month of record within square, not the locality within square

