

# African Marsh Owl

*Asio capensis*

170 plots from 238 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

	Seasonality		
	presence	bs	eggs
January	32	1	2
February	23	-	-
March	17	-	-
April	25	1	1
May	23	1	1
June	8	2	-
July	16	-	-
August	15	-	1
September	12	-	-
October	16	-	-
November	18	-	-
December	21	-	-
month 13	2	-	-


recorded from  
**61 squares**  
17%

**Status**

- Presence
- Breeding season
- Month with eggs

from Britton (1980)

The nominate race is a locally common resident in open grasslands, moorlands, marshes and swampy hollows up to 3000 m. In Kenya it is fairly common and widespread in the rift valley and adjoining plateaux north to Lake Turkana at Kalakol, and ranging rather uncommonly west to Trans Nzoia and Nandi, and east to Aruba in Tsavo East NP. In Tanzania it ranges from Arusha NP, Tarangire NP, Lake Manyara, Serengeti NP and Mwanza south to the Rukwa Valley, east to the lowlands of Kilimanjaro and the Tanga hinterland. In Uganda it is a rather uncommon bird of the west, ranging north and east to Kabalega Falls NP and Mengo.



**Tanzania Bird Atlas**  
Preliminary Map  
July 2015  
From 1,135,191  
Database records

J F M A  
M J J A  
S O N D  
# # # #

symbols give month of  
record within square, not  
the locality within square