


# African Marsh Owl

*Asio capensis*

181 plots from 260 records



recorded from  
63 squares  
18 %

Virgin Yellow-vented Bulbul squares **4**  
Single YvB month **13**  
Two YvB months **18**

- Presence
- ▲ Breeding season
- Month with eggs

from Britton (1980)  
The nominate race is a locally common resident in open grasslands, moorlands, marshes and swampy hollows up to 3000 m. In Kenya it is fairly common and widespread in the rift valley and adjoining plateaux north to Lake Turkana at Kalakol, and ranging rather uncommonly west to Trans Nzola and Nandi, and east to Aruba in Tsavo East NP. In TZ it ranges from Arusha NP Tarangire NP, Lake Manyara, Serengeti NP and Mwanza south to the Rukwa Valley, east to the lowlands of Kilimanjaro and the Tanga hinterland. In UG it is a rather uncommon bird of the west, ranging north and east to Kabalega Falls NP and Mengo.

	Seasonality	
	presence	bs eggs
January	34	1 2
February	25	- -
March	18	- -
April	25	1 1
May	23	1 1
June	9	2 1
July	18	1 -
August	18	- 1
September	13	- -
October	17	- -
November	22	- -
December	23	- -
month 13	2	



Tanzania Bird Atlas  
Preliminary Map  
March 2017  
From 1,180,093  
Database records

J F M A  
M J J A  
S O N D  
#####  
symbols give month of  
record within square, not  
the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E