

Eurasian Nightjar

Caprimulgus europaeus

108 plots from 169 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Status

- Presence
- Breeding season
- Month with eggs


Seasonality

	presence	bs	eggs
January	29	-	-
February	33	-	-
March	56	-	-
April	25	-	-
May	1	-	-
June	-	-	-
July	-	-	-
August	-	-	-
September	2	-	-
October	10	-	-
November	24	-	-
December	13	-	-

from Britton (1980).

A fairly common and widespread Palearctic migrant in a variety of bushed and wooded habitats and cultivation from sea-level to 3000 m, mainly on passage during October-November and March-April. There are records of the nominate race and *unwini* from scattered localities throughout East Africa, and most specimens from E Tanzania are *unwini*.

Some birds referred to the nominate race might be the slightly smaller *meridionalis*, which has been recorded from C Kenya and interior N Tz, though it is doubtfully separable in Africa. About 30 per cent of birds handled at Ngulia in E Ke in November-December are referable to either *unwini* or *plumipes*.



recorded from
62 squares
17 %



Tanzania Bird Atlas
Preliminary Map
March 2015
From 1,120,060
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square