

Blue-naped Mousebird

Urocolius macrourus

1,023 plots from 3,371 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality

	presence	bs	eggs
January	370	2	2
February	319	1	-
March	330	1	-
April	209	1	-
May	191	2	1
June	251	1	-
July	338	-	-
August	303	1	1
September	241	2	-
October	269	5	-
November	212	1	-
December	314	2	-

It would be really interesting to confirm that both this species and Red-faced occur around Kilwa. Photos please.


from Britton (1980)

A common resident of bushland open woodland and bushed grassland typically in arid and semi-arid areas below 1600 m. Four races occur *abyssinicus* along the northern border of Kenya east to Moyale *griseogularis* throughout Uganda except Karamoja, south to Lake Urigi in NW Tanzania *massaicus* in NE Tanzania from Dar es Salaam and Dodoma to Shinyanga and the lowlands of Arusha and Kilimanjaro *pulcher* is widespread in Tanzania and Kenya north of *massaicus*, west to Karamoja and Ukerewe Island, and south of *massaicus* from Morogoro to Lakes Rukwa and Malawi, allopatric with *U. indicus* with which it forms a superspecies.

recorded from
173 squares
49 %

Status

- Presence
- Breeding season
- Month with eggs



Tanzania Bird Atlas
Preliminary Map
July 2015
From 1,135,191
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square