

Eurasian Bee-eater

Merops apiaster

1,073 plots from 3,724 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Status
 ● Presence
 ● Breeding season
 ● Month with eggs


Seasonality
 presence bs eggs

Month	presence	bs	eggs
January	486	-	-
February	492	-	-
March	600	-	-
April	226	-	-
May	28	-	-
June	4	-	-
July	2	-	-
August	16	-	-
September	281	-	-
October	710	-	-
November	519	-	-
December	346	-	-
month 13	14	-	-

recorded from
 280 squares
 80 %

from Britton (1980).
 An often numerous Palearctic visitor which occurs throughout East Africa on passage, September-November and late March-early May, exceptionally in July and August. Parties frequent open country with scattered trees, woodland and forest edges from sea-level to 3000 m. Smaller numbers winter in most parts of Tanzania, north to W and C Kenya and SW Uganda.

Clearly this species really does occur throughout the country, the gaps shown here relate more to absence of observers than lack of birds.



Tanzania Bird Atlas
 Preliminary Map
 March 2015
 From 1,120,060
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of
 record within square, not
 the locality within square