


Trumpeter Hornbill

Bycanistes bucinator

665 plots from 1,723 records



Virgin Yellow-vented Bulbul squares **4**

Single YvB month **13**

Two YvB months **18**

Seasonality
presence by eggs

| Month | presence | by | eggs |
|-----------|----------|----|------|
| January | 157 | 1 | - |
| February | 112 | 2 | - |
| March | 146 | - | - |
| April | 90 | - | - |
| May | 90 | - | - |
| June | 159 | - | - |
| July | 180 | 2 | 1 |
| August | 169 | - | - |
| September | 180 | 1 | 2 |
| October | 161 | - | 5 |
| November | 145 | - | - |
| December | 124 | - | - |

from Britton (1980)
A common and sometimes abundant resident and wanderer in forest and woodland up to 1500 m with a curious patchy distribution. While ranging south along the coast from Vwitu and the Tana River to Mikindani, it is unrecorded in coastal KE south of Sokoke F. and uncommon or absent in much of coastal TZ. There are scattered records from much of interior TZ south of 5° S, west to Lake Tanganyika and north to the Usambaras, the foothills of the N. Pare Mts and Mt Kilimanjaro, and the Kahambwe River in Kibondo. In KE it ranges west of Kiboko, Kitui, Thika, Embu and Meru, mainly south of the equator.

I've long been uncomfortable with the statement in Britton that this species is "uncommon or absent in much of coastal Tanzania". Good to correct this misconception and a clear reminder of how poorly known SE Tanzania was until quite recently.



Tanzania Bird Atlas
Preliminary Map
February 2017
From 1,173,386
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E