

Southern Ground Hornbill

Bucorvus leadbeateri


1,082 plots from 2,890 records

Virgin Yellow-vented Bulbul squares **7**
 Single YvB month **14**
 Two YvB months **20**

	Seasonality		
	presence	bs	eggs
January	295	3	3
February	223	4	1
March	221	3	1
April	188	2	1
May	143	3	-
June	251	1	1
July	294	-	-
August	287	-	1
September	246	1	-
October	259	1	-
November	229	2	1
December	203	4	3
month 13	15		

recorded from
253 squares
72 %

Status
 ● Presence
 ● Breeding season
 ● Month with eggs



Tanzania Bird Atlas
 Preliminary Map
 June 2015
 From 1,130,669
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of
 record within square, not
 the locality within square

from Britton (1980)
 A widespread and sometimes common resident and wanderer from sea-level to 3000 m, typically in less arid country than *abyssinicus*. There are records from most parts of Tz though few from West Lake or the coast. In Ke it ranges north to Witu, Tsavo NP (E and W), Embu, Aberdares, Eldama Ravine, Lessos, Eldoret, Nandi Hills, the S. Nandi Forest, Kipsait and the Turkwell River. There are scattered records from the north side of L. Victoria west to near Entebbe, and it is reasonably common in South Nyanza (north to Ahero) in W Ke, but it is no more than a vagrant in lowland areas in between.