

# Green Barbet

*Stactolaema olivacea*

115 plots from 367 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**


	Seasonality		
	presence	bs	eggs
January	14	2	-
February	33	1	2
March	26	1	-
April	23	-	-
May	13	-	-
June	20	2	-
July	50	1	-
August	53	-	-
September	37	7	-
October	14	1	1
November	29	7	-
December	28	-	-
month 13	2		

from Britton (1980). Typically a bird of forests from sea-level to 1800 m, but common in both wooded and forested habitats at Sokoke in coastal Kenya. The nominate race is confined to the coastal fringe in Kenya, from Shimoni to the Tana River, including Rabai, Ribe and the Shimba Hills, while in E Tanzania it is a bird of the highlands, from the Usambaras (East and West) to the Ngurus and Mahenge. Three other races occur in E Tanzania: *rungweensis* in the highlands at the north end of Lake Malawi, from Umalila and Isoko to Mt Rungwe and the Poroto Mts; *woodwardi* at Nchingidi on the Rondo Plateau; *uluguruensis*, described by Ripley and Heinrich (1969), in the Ulugurus.

recorded from 29 squares 8.2%

### Status

- Presence
- Breeding season
- Month with eggs



Tanzania Bird Atlas  
Preliminary Map  
May 2014  
From 1,070,127  
Database records

J F M A  
M J J A  
S O N D  
# # # #

symbols give month of record within square, not the locality within square

Birds of Africa vol III page 427 do not accept *uluguruensis* which they include with nominate birds. They do include *howelli* described by Jensen and Stuart in the 1980s from the East Udzungwa Mts.

BoA mention the apparent gap between our *woodwardi* in the SE and what they consider the same race known from the Ngoye forest in KwaZulu Natal. With so much fieldwork in Mozambique this past decade or so does anyone know if this gap still exists or was it simply a knowledge gap.