

Red and Yellow Barbet

Trachyphonus erythrocephalus

269 plots from 1,069 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

	Seasonality		
	presence	bs	eggs
January	126	3	1
February	92	-	-
March	82	2	-
April	73	1	-
May	65	3	-
June	71	2	-
July	102	-	-
August	94	-	-
September	85	2	-
October	87	1	-
November	75	1	1
December	90	8	1
mothn 13	1		

recorded from
64 squares
18 %

Status
 ● Presence
 ● Breeding season
 ● Month with eggs

Some serious work for us here as we attempt to establish why there are so many records well away from the obvious core habitat. Does this bird MOVE? Most will surely be ID and entry errors but note that Britton also included Mikumi NP in the range of this species.


from Britton (1980)
 The habitat requirements and ranges of this and the previous species are broadly similar. The two differ greatly in size, and co-exist on the same ground in many areas, though this species is often in more open areas than *darnaudii*. The race *versicolor* is wide-ranging in N Ke south to the Kerio Valley, L. Bogoria and Meru NP, extending to NE Ug from Kidepo Valley NP to Moroto, and possibly in northern border areas to the west. The nominate race occurs in S Ke and NE Tz, including Mara GR, the Ngong area, the Tsavo region, the lower Tana River, Lake Manyara and Mikumi NP.



Tanzania Bird Atlas
 Preliminary Map
 May 2015
 From 1,128,849
 Database records

J F M A
 M J J A
 S O N D
 # # # #

symbols give month of record within square, not the locality within square



29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E