

Usambiro Barbet

Trachyphonus usambiro

184 plots from 656 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	83	1	1
February	81	3	1
March	73	2	-
April	43	2	-
May	27	1	1
June	54	-	-
July	73	-	-
August	45	-	-
September	25	-	-
October	45	-	-
November	45	2	-
December	46	2	-

recorded from
37 squares
10.6 %

Status

- Presence
- Breeding season
- Month with eggs

These records in the Natron Rift will be double checked as will the odd record well away from the core range. It should be d'Arnaud's in the dry country south of the Mbulu Plateau.

from Britton (1980)
This East African endemic is widespread and common in bushed and wooded grassland and open bushland in semi-arid country south and southeast of Lake Victoria at 1100-2100 m. It ranges from Naivasha, Morijo and Aneko in SW Kenya to Serengeti, Hanang, Wembere and Mwanza (west to Ukiriguru and Usambiro) in N Tanzania.



Tanzania Bird Atlas
Preliminary Map
April 2015
From 1,128,849
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

