

Red-throated Wryneck

Jynx ruficollis

40 plots from 57 records

Virgin Yellow-vented Bulbul squares **9**


Single YvB month **16**

Two YvB months **24**

Status
 ● Presence
 ● Breeding season
 ● Month with eggs

Seasonality	presence	bs	eggs
January	3	-	-
February	7	-	-
March	1	-	-
April	6	-	-
May	4	-	-
June	5	-	-
July	4	-	-
August	3	-	-
September	3	1	-
October	4	-	-
November	7	-	-
December	7	1	1

recorded from
20 squares
5.7%



from Britton (1980). The nominate race occurs in woodlands, open areas with trees and forest edges at 600-3000 m (seldom below 1700 m in Kenya). While scarce and local over most of its East African range it is quite common in parts of Kenya and NW Tanzania. In Kenya it ranges from Mt Elgon, the Wei-Wei River, Maralal, the Aberdares and Murang'a to Rapogi, Mara GR, the Nguruman Hills, Nairobi and Simba, ranging north of Elgon to Sebei (Uganda). Elsewhere in Uganda it is known from scattered localities in the west, from Kigezi to Masaka, Mubende and Bugoma. In Tanzania it occurs in West Lake and Kibondo in the northwest and in the Ruvuma Region east to Nakapanya in the south. A bird collected at Kingolwira in E Tanzania regarded as a possible hybrid between this and the next species by Desfayes (1969), is best regarded as a variant of *ruficollis* (Short and Bock 1972).



Tanzania Bird Atlas
 Preliminary Map
 January 2014
 From 1,056,327
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of
 record within square, not
 the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E