

# Cardinal Woodpecker

*Dendropicos fuscescens*

1,248 plots from 3,005 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**


Seasonality  
presence bs eggs

Month	presence	bs	eggs
January	272	2	1
February	229	1	-
March	247	1	-
April	185	3	1
May	174	-	-
June	255	2	1
July	297	6	1
August	295	8	1
September	241	2	2
October	241	3	-
November	217	9	3
December	277	9	1
month 13	18		

recorded from  
271 squares  
77 %

### Status

- Presence
- Breeding season
- Month with eggs



Tanzania Bird Atlas  
Preliminary Map  
April 2015  
From 1,128,849  
Database records

J F M A  
M J J A  
S O N D  
# # # #

symbols give month of  
record within square, not  
the locality within square

from Britton (1980)  
A wide-ranging and common bird of open woodland, wooded grassland and bushland up to 2600 m. Four races occur *hemprichii* in N and E Ke, west and south to Moyale, Marsabit, the N Uaso Nyiro and Tana rivers, and the coast from Lamu to Sokoke Forest, extending to the lowlands of Kilimanjaro in NE Tz and Kidepo Valley NP in NE Ug *massaicus* elsewhere in Ke north and east of the highlands, south to Oldeani, Babati and Moshi in NE Tz, and in NE Ug from Teso to Moroto *hartlaubii* in most of Tz, inc Zanzibar, north to the Chyulus in SE Ke *lepidus* in the remaining areas of Ke and Ug south to Bukoba in NW Tz, and localized in the E Usambaras in NE Tz (at 500-1150 m). The last is more of a forest bird than the other races, and has been regarded as specifically distinct (in the polytypic *D. lafresnayi*) by Ripley and Heinrich (1969).