

Cardinal Woodpecker

Dendropicus fuscescens

1,206 plots from 2,820 records

○ hemprichii ○ lepidus
○ massaicus ○ lafresnayi
○ hartlaubii

ONLY 7 NAMED LOCALITIES IN BRITTON


recorded from
268 squares
76%

Status

- Presence
- Breeding season
- Month with eggs

Seasonality

	presence	bs	eggs
January	267	2	1
February	220	1	-
March	226	1	-
April	176	3	1
May	162	-	-
June	207	2	1
July	273	6	1
August	280	8	1
September	237	2	2
October	229	3	-
November	203	8	3
December	266	9	1
month 13	18		



from Britton (1980).
 A wide-ranging and common bird of open woodland, wooded grassland and bushland up to 2600 m. Four races occur *hemprichii* in N and E Kenya, west and south to Moyale Marsabit, the N Uaso Nyiro and Tana rivers, and the coast from Lamu to Sokoke Forest, extending to the lowlands of Kilimanjaro in NE Tanzania and Kidepo Valley NP in NE Uganda *massaicus* elsewhere in Kenya north and east of the highlands, south to Oldeani, Babati and Moshi in NE Tanzania, and in NE Uganda from Teso to Moroto *hartlaubii* in most of Tanzania, including Zanzibar, north to the Chyulus in SE Kenya. *lepidus* in the remaining areas of Kenya and Uganda, south to Bukoba in NW Tanzania, and localized in the East Usambaras in NE Tanzania (at 500-1150 m) The last is more of a forest bird than the other races, and has been regarded as specifically distinct (in the polytypic *D. lafresnayi*) by Ripley and Heinrich (1969).



Tanzania
BIRD ATLAS

Tanzania Bird Atlas
Preliminary Map
March 2014
From 1,061,639
Database records

J F M A
M J J A
S O N D
#

centralis in northern Zambia

hartlaubii in northern Malawi

symbols give month of record within square, not the locality within square