

Foxy Lark

Calendulauda alopex

379 plots from 1,099 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	107	2	-
February	100	2	-
March	107	8	-
April	91	4	2
May	67	4	-
June	72	2	-
July	95	-	-
August	85	1	-
September	59	-	-
October	80	1	-
November	81	4	-
December	122	3	-

recorded from
80 squares
23 %

Status
● Presence
● Breeding season
● Month with eggs

from Britton (1980)
The race *intercedens* is a locally common resident of arid or semi-arid country at 500-1800 m, frequenting open and bushed grasslands on dry sandy soils, the colour of which results in distinct colour variations in plumage. In interior Kenya it ranges widely, mainly in the rift valley or eastwards, from Lodwar, Pokot and Marsabit to the Loita Plains, Amboseli, the Tsavo region and Taru, extending to Karamoja in NE Uganda and to N Tz from Serengeti to Dodoma, Tarangire NP and the foothills of the North Pare Mts.



Tanzania Bird Atlas
Preliminary Map
May 2015
From 1,130,669
Database records

J F M A
M J J A
S O N D

symbols give month of
record within square, not
the locality within square

