



## Fork-tailed Drongo *Dicrurus adsimilis*

2277 plots from 10,010 records

Virgin Yellow-vented Bulbul squares **4**  
 Single YvB month **13**  
 Two YvB months **18**

### Seasonality presence vs eggs

Month	presence	vs	eggs
January	827	18	4
February	754	12	4
March	776	6	4
April	566	9	-
May	615	10	4
June	966	6	1
July	1,225	4	-
August	986	2	1
September	823	-	3
October	816	8	4
November	722	19	10
December	759	18	10
month 15			16

from Britton (1980).  
 Typically a widespread and common resident of forest edge, dry woodland, wooded grassland and cultivation up to 2200 m, though the race *coracinus* is a forest bird. The two non-forest races occupy drier habitats throughout East Africa, including Zanzibar, being particularly prominent in acacia country. The northern race *divaricatus* intergrades with nominate birds around the equator.

**HAS ANYONE EVEN SUSPECTED MOVEMENT IN THIS BIRD**



Tanzania Bird Atlas  
 Preliminary Map  
 December 2016  
 From 1,173,239  
 Database records

J F M A  
 M J J A  
 S O N D  
 ■ ■ ■ ■

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E