


Lesser Swamp Warbler

Acrocephalus gracilirostris


324 plots from 684 records

Virgin Yellow-vented Bulbul squares	9
Single YvB month	12
Two YvB months	21

	Seasonality		
	presence	bs	eggs
January	74	3	-
February	72	-	-
March	85	2	-
April	42	3	-
May	45	1	1
June	42	-	3
July	41	1	1
August	45	-	-
September	53	-	-
October	46	-	-
November	63	-	-
December	57	2	1
month 13	1		



from Britton (1980).
 Wide-ranging up to 2400 m, though absent from much of NW and SE Tanzania, N and E Kenya and N Uganda. The race *jacksoni* is abundant in low shrubbery on rocky islands and similar mainland sites on the shores of Lake Victoria in Uganda and W Kenya, but local elsewhere in the Lake Victoria basin and W Uganda, mainly in swamp edges, from Butiaba and Rwenzori NP to Tororo, Ukwala, Kisumu and Waturi. Other races are common birds of Typha and papyrus swamps *leptorhyncha* in the coastal lowlands from Dar es Salaam and Soga to the lower Tana River, including Zanzibar and Pemba, inland to Simba, Kibwezi, Taveta, the East and West Usambaras, the Kilombero River, Iringa and Songea *parvus* in the Kenya highlands from Eldoret, Kapsabet and Molo to Laikipia and Nairobi, and in N and W Tanzania from Mt Kilimanjaro and Monduli to Mbulu, Kome Island, Tabora and the shores of Lake Tanganyika.



Tanzania Bird Atlas
 Preliminary Map
 February 2015
 From 1,113,718
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square