

Grey Wren Warbler

Camaroptera simplex

213 plots from 518 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Status

- Presence
- Breeding season
- Month with eggs


Seasonality

	presence	bs	eggs
January	56	-	-
February	34	-	-
March	37	2	1
April	37	6	1
May	45	1	-
June	30	-	-
July	52	-	-
August	49	-	-
September	26	-	-
October	21	-	-
November	38	-	-
December	80	1	-
month 13		1	

recorded from
37 squares
10.5%

C. s. undosa is now treated as a full species. Pale Wren Warbler in some lists, Miombo Wren Warbler in others.

from Britton (1980). Wide-ranging and reasonably common, represented by two well-marked races which might be better regarded as separate species. The race *undosa* is typically associated with thin cover on termite mounds in miombo woodland, but also occurs in the tangled undergrowth of bushland and woodland, at 900—1700 m. It ranges throughout W Tanzania from Ufipa and Mbeya to N Tabora, Kibondo, Biharamulo and Ngara, locally east to Ruaha NP, Sao Hill, the northern edge of the Iringa highlands, Bahi and the foothills of the Ulugurus, and north to the Mara River and Loita Hills in SW Kenya. The nominate race occurs in bushland and woodland in arid and semi-arid country at 200—1300 m, usually in tangled undergrowth and often in the vicinity of seasonal watercourses. It ranges widely north, east and south of the Kenya highlands, extending to Moroto in NE Uganda and border areas of NE Tanzania from Lake Natron and Loliondo to the Pangani River. This and the next species are members of a superspecies. In areas of apparent sympatry in E Tanzania, *stierlingi* is found at lower elevations.



Tanzania Bird Atlas
Preliminary Map
January 2014
From 1,056,327
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square