

Silverbird


Empidonis semipartitus

348 plots from 1,614 records

Virgin Yellow-vented Bulbul squares	5
Single YvB month	12
Two YvB months	18

Seasonality

	presence	bs	eggs
January	178	5	2
February	155	6	2
March	111	5	1
April	57	2	-
May	152	2	2
June	192	2	3
July	206	1	-
August	110	-	-
September	70	-	-
October	99	1	-
November	88	2	2
December	150	6	2



Note this isolated population inhabiting Acacia floodplain in a sea of Miombo woodland.

recorded from 69 squares 19.7%

Status
 ● Presence
 ● Breeding season
 ● Month with eggs



Tanzania Bird Atlas
 Preliminary Map
 September 2016
 From 1,165,026
 Database records

J F M A
 M J J A
 S O N D
 # # # #

symbols give month of record within square, not the locality within square

from Britton (1980).
 Locally common in bushed grassland (especially acacia) in moderately arid areas at 400-1500m in N Uganda, NW Kenya, SW Kenya and interior N Tanzania. In the north it occurs from West Nile to S Turkana, south to Bunyoro, Buruli, Sebei, Kongelai, Lake Baringo and Isiolo. An apparently separate population occurs along the east shore of Lake Victoria from Yimbo to Mwanza, thence south to interior Tanzania to Shinyanga and C Tabora, east to Mara GR in Kenya and W Arusha and N Dodoma in Tanzania. Its absence from high rainfall areas between Kongelai and Yimbo is apparently real, yet it is common at Rapogi and in other equally verdant parts of South Nyanza. Most are probably resident, but there are records of apparent wanderers from as far afield as Central Island in Lake Turkana and Same in NE Tanzania, as well as occasional birds in the Kenya highlands (up to 1800m) at Nairobi, Lake Nakuru and Rongai.