

# Striped Pipit

*Anthus lineiventris*

110 plots from 158 records

Virgin Yellow-vented Bulbul squares **4**

Single YvB month **13**

Two YvB months **18**

Seasonality  
presence by eggs

January	13	-	-
February	12	-	-
March	16	-	-
April	7	-	-
May	5	-	-
June	7	-	-
July	12	-	1
August	14	-	-
September	16	-	-
October	16	-	-
November	19	2	-
December	16	1	1

Rarely an easy bird, there are even month gaps for Iringa where it occurs on the fringes of the town in peoples gardens. Do note that it is surely resident on the lower slopes of the West Usambara above Mombo but so few records ??

recorded from  
60 squares  
17 %

We must assume that when people go birding out west they dont climb the hills. Recent records from Uganda suggest it is far more common in W TZ than suggested here or in Britton (1980).

- Presence
- Breeding season
- Month with eggs

from Britton (1980).  
A locally distributed and generally scarce resident of rocky wooded hillsides in E Tanzania and SE Kenya, mainly in the highlands at 1000-2000 m, ranging from Songea and Tunduru to the North Pare Mts, Mt Kilimanjaro, the Mbulu Escarpment and the Taita and Chyulu hills. One in miombo woodland on a rocky hill south of Kibondo on 21 March 1961 is the only record for W Tanzania.

J F M A  
M J J A  
S O N D  
# # # #

symbols give month of record within square, not the locality within square



Tanzania Bird Atlas  
Preliminary Map  
March 2017  
From 1,180,093  
Database records

