

Palm-nut Vulture

Gypohierax angolensis

627 plots from 1,770 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Status


- Presence
- Breeding season
- Month with eggs

Seasonality

	presence	bs	eggs
January	189	-	-
February	123	6	-
March	141	3	-
April	92	1	-
May	90	5	-
June	145	4	-
July	161	5	1
August	160	2	3
September	174	5	3
October	154	1	1
November	148	8	1
December	144	-	-

As with all raptors proving eggs in nest is far from easy but do look out for young in nest and do try and age them and work backwards to at least egg months.

recorded from 141 squares 40%



from Britton (1980). Resident in lowland areas with *Raphia* or *Elaeis* palms, mainly in W Tanzania, S and W Uganda, and the coastal lowlands between the Rufiji River and Witu, including Zanzibar and Pemba. It is common along the shores of Lake Tanganyika and locally at the coast, especially on Pemba and the neighbouring mainland, inland to Amani and north to the Shimba Hills. Apart from Trans Nzoia and Mweiga in the Kenya highlands, all other recorded sites are below 1400 m, including Ruaha NP, Kidepo Valley NP, Tsavo West NP (at Lake Jipe and Mzima Springs), Lake Manyara, the foothills of the Ulugurus, Kibwezi, the Tana River upstream to Meru NP, and at springs and streams around Mt Kilimanjaro.

Tanzania Bird Atlas
Preliminary Map
February 2014
From 1,061,639
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E